

JT EATON™

NECTUS™

SOFT BAIT

2G SECOND GENERATION
RODENTICIDE™

-PEANUT BUTTER FLAVOR-

ACTIVE INGREDIENT:

Bromadiolone
(CAS #28772-56-7):..... 0.005%

OTHER INGREDIENTS*:..... **99.995%**

TOTAL.....100.000%

*Contains Denatonium Benzoate

ACCEPTED
FOR REGISTRATION

JULY 20 2020

New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration

Doc ID: 568588

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

SEE BACK PANEL FOR ADDITIONAL
PRECAUTIONARY LABELING.

JT EATON™
SINCE 1932

Manufactured in Slovenia for: J.T. EATON & CO., INC.
1393 East Highland Road Twinsburg, OH 44087
U.S.A. © 2020 | www.jteaton.com

JT EATON™, 2G Second Generation Rodenticide™ and
Nectus™ are trademarks of J.T. EATON & CO., INC.

**KILLS
RATS AND MICE**
KILLS WARFARIN RESISTANT
NORWAY RATS

NET CONTENTS:

16lbs. (7.3 kgs)

EPA Reg. No. 82744-2-56
EPA Est. No. 82744-SLO-001

ITEM# 716-S

FIRST AID

HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE.

IF SWALLOWED:	<ul style="list-style-type: none">• Call a poison control center, doctor, or 1-800-222-1222 immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center, doctor, or 1-800-222-1222 immediately for treatment advice.
IF IN THE EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center, doctor, or 1-800-222-1222 immediately for treatment advice.
NOTE TO PHYSICIAN: If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.	

TREATMENT FOR PET POISONING: If animal eats bait, call veterinarian at once.

NOTE TO VETERINARIAN: Anticoagulant Bromadiolone: For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K1. If needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through the skin. Keep away from children, domestic animals and pets. Causes moderate eye irritation. Do not get in eyes, on skin, or on clothing. All handlers (including applicators) must wear shoes plus socks, and waterproof gloves. Any person who retrieves carcasses or unused bait following application of this product must wear waterproof gloves.

USER SAFETY REQUIREMENTS: Follow manufacturer's instruction for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to mammals and birds. Dogs, cats, and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. This pesticide is toxic to fish. Do not apply this product directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may also be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash water or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

IMPORTANT: Do not expose children, pets, or nontarget animals to rodenticides. To help to prevent accidents:

1. Store unused product out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.

3. Dispose of product container and unused, spoiled, or unconsumed bait as specified on this label. Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used if children, pets, non-target mammals, or birds may access the bait.

USE RESTRICTIONS: This product can only be used to control Norway rats, roof rats, and house mice in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port of terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting, beyond 100 feet from a structure as defined above, is prohibited. This product must not be applied directly to food or feed crops. Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food. When used in USDA inspected facilities, this product must be applied in tamper-resistant bait stations. Burrow baiting with JT EATON™ NECTUS™ SOFT BAIT 2G Second Generation Rodenticide™ is prohibited.

DO NOT SELL THIS PRODUCT IN INDIVIDUAL CONTAINERS HOLDING LESS THAN 16 POUNDS OF BAIT.

SELECTION OF TREATMENT AREAS: After removing as much food, water and harborage as possible, determine dry areas where rats and mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, beside burrows, in corners and concealed places, between floors and walls, or in locations where signs of rats or house mice have been seen. Protect bait from rain and snow.

APPLICATION DIRECTIONS: Each bag of JT EATON™ NECTUS™ SOFT BAIT 2G Second Generation Rodenticide™ contains approximately 0.38 ounces (10.75 grams) of bait.

Norway Rats and Roof Rats: Apply 10 to 42 bait bags per placement for controlling rats. Space placements 15 to 30 feet apart at appropriate locations within the infested area. Maintain a continuous supply of fresh bait for at least 10 days or until signs of new feeding by rats on bait cease to appear.

House Mice: Apply 1 bait bag per placement, usually spaced 8 to 12 feet apart. Two bags may be needed at points of very high mouse activity. Maintain a constant supply of fresh bait for 15 days or until signs of mouse activity cease.

Follow-up: While wearing gloves, replace contaminated or spoiled bait immediately. Collect and properly dispose of dead animals and leftover bait and bait bags, as well as any bait that has become accessible to non-target animals. To discourage re-infestation, limit sources of rodent food, water, and harborage as much as possible. If re-infestation does occur, repeat treatment. For continuous sources of infestation, set up permanent bait stations and replenish them as needed.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.

PESTICIDE STORAGE: Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be placed in trash or delivered to an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container.

Triple rinse (or equivalent), then offer for recycling or reconditioning; or puncture and dispose of in sanitary landfill; or by incineration. In most states, burning is not allowed. Remove all plastic bags after the treatment and dispose of them in accordance with local requirements.

WARRANTY: To the extent consistent with applicable law, seller makes no warranty expressed or implied, concerning the use of this product other than indicated on the label. To the extent consistent with applicable law, buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

FNW716SB2JTD
EPA03272020/INV04172020

